



**Dr. Joe Ables, PA**  
499 CR 3198  
Decatur, TX 76234  
940-389-1738

**Deer Vaccinations:**

- A. Covexin 8- 2.5 cc SQ**
- B. Virashield 6 + HS (Haemophilus somnus) – 2.5 cc SQ**
- C. Prespense HM – 1 cc SQ**
- D. Fusogard – 1 cc SQ**
- E. Staybred VL5– 1 cc SQ**

*\*\*\* I recommend mixing A and B in one syringe for a total of 5 cc and use a 16 gauge needle, left side of the neck. I recommend mixing C, D, and E in one syringe and giving a total of 3 cc SQ on the right side of the neck. Changing sides will distribute out soreness. SQ = subcutaneous (under the skin)*

**Antibiotics / Dewormers:**

- A. Zactran – 2cc per 100 lbs SQ (lasts about 10 days)**
- B. Excede – 2 cc per 100 lbs SQ (lasts about 7 days), if sick, give 3 cc per 100 lbs SQ**
- C. Banamine (flunixin meglumine/ “PreVail”) – 1 cc per 100 lbs IM one time only**
- D. Dectomax- 2cc per 100 lbs IM**

*\*\*\* All long acting medications (antibiotics and dewormers) should be give SQ (subcutaneous) / under the skin*

*\*\*\* Body weights will not affect vaccine dosages of adults and fawns. However, body weights will affect antibiotic and dewormer dosages between adults and fawns*

**Glossary:**

SQ- subcutaneous- under the skin

IM- in the muscle

IV- in the vein