



**Dr. Joe Ables, PA**  
499 CR 3198  
Decatur, TX 76234  
940-389-1738

**Spring Deer Vaccinations:**

- A. Covexin 8- 2.5 cc SQ
- B. Virashield 6 + HS (Haemophilus somnus) – 2.5 cc SQ
- C. Prespense HM – 1 cc SQ
- D. Fusogard – 1 cc SQ
- E. Clostridium A – 1 cc SQ

*\*\*\* I recommend mixing A and B in one syringe for a total of 5 cc and use a 16 gauge needle, left side of the neck. I recommend mixing C, D, and E in one syringe and giving a total of 3 cc SQ on the right side of the neck. Changing sides will distribute out soreness. SQ = subcutaneous (under the skin)*

**Antibiotics / Dewormers:**

- A. Zactran – 2cc per 100 lbs SQ (lasts about 10 days)
- B. Excede – 2 cc per 100 lbs SQ (lasts about 7 days), if sick, give 3 cc per 100 lbs SQ
- C. Banamine (flunixin meglumine/ “PreVail”) – 1 cc per 100 lbs IM one time only
- D. Dectomax 10% - 2 cc per 100 lbs IM
- E. Ivomec plus- 2 cc per 100 lbs SQ- best for areas (low water areas) with liver fluke problems.

**Safe for pregnant doe. In the fall, you can drench with Valbazen.**

*\*\*\* All long acting medications (antibiotics and dewormers) should be give SQ (subcutaneous) / under the skin\*\*\* Body weights will not affect vaccine dosages of adults and fawns. However, body weights will affect antibiotic and dewormer dosages between adults and fawns*

**Glossary:**

SQ- subcutaneous- under the skin

IM- in the muscle, IV- in the vein